

*11th Annual Strawberry Production Meeting in Ventura County
Camarillo, August 31, 2012*

Lygus, Thrips, Corn Earworm and Drosophila

Frank Zalom

Department of Entomology
University of California, Davis

Lygus Control Update - Insecticides 2011



Registered and Candidate Insecticides

Pesticide	Chemical	Subgroup	Target Site of Activity	IRAC #
Lannate	methomyl	carbamate	acetylcholine esterase inhibitor	1A
Malathion	malathion	organophosphate	acetylcholine esterase inhibitor	1B
Dibrom	naled	organophosphate	acetylcholine esterase inhibitor	1B
Brigade	bifenthrin	pyrethroid	sodium channel modulator	3
Danitol	fenpropathrin	pyrethroid	sodium channel modulator	3
Actara	thiamethoxam	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Assail	acetamiprid	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Rimon	novaluron	benzoylureas	inhibitor of chitin biosynthesis, type 0	15
Prevam	borax	borax	nonspecific inhibitor	uncl
Belay	clothianidin	neonicotinoid	nicotinic Ach receptor agonist/ antagonist	4A
Beleaf	flonicamid	flonicamid	nonspecific feeding blocker	9C
Closer	sulfoxaflor	sulfilimine	--	uncl
--	tolyfenpyrad	METI insecticide	Mitochondrial complex I electron transport inhibitor	21A

Not registered

Registered

Not registered for use on strawberries, but under study

Nerve Poisons

Pesticide	Chemical	Subgroup	Target Site of Activity	IRAC #
Malathion	malathion	organophosphate	acetylcholine esterase inhibitor	1B
Dibrom	naled	organophosphate	acetylcholine esterase inhibitor	1B
Brigade	bifenthrin	pyrethroid	sodium channel modulator	3
Danitol	fenpropathrin	pyrethroid	sodium channel modulator	3
Actara	thiamethoxam	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Assail	acetamiprid	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Rimon	novaluron	benzoylureas	inhibitor of chitin biosynthesis, type 0	15
Prevam	borax	borax	nonspecific inhibitor	uncl
Belay	clothianidin	neonicotinoid	nicotinic Ach receptor agonist/ antagonist	4A
Beleaf	flonicamid	flonicamid	nonspecific feeding blocker	9C
Closer	sulfoxaflor	sulfilimine	--	uncl
--	tolyfenpyrad	METI insecticide	Mitochondrial complex I electron transport inhibitor	21A

Insect Growth Regulator

Pesticide	Chemical	Subgroup	Target Site of Activity	IRAC #
Malathion	malathion	organophosphate	acetylcholine esterase inhibitor	1B
Dibrom	naled	organophosphate	acetylcholine esterase inhibitor	1B
Brigade	bifenthrin	pyrethroid	sodium channel modulator	3
Danitol	fenpropathrin	pyrethroid	sodium channel modulator	3
Actara	thiamethoxam	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Assail	acetamiprid	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Rimon	novaluron	benzoylureas	inhibitor of chitin biosynthesis, type 0	15
Prevam	borax	borax	nonspecific inhibitor	uncl
Belay	clothianidin	neonicotinoid	nicotinic Ach receptor agonist/ antagonist	4A
Beleaf	flonicamid	flonicamid	nonspecific feeding blocker	9C
Closer	sulfoxaflor	sulfilimine	--	uncl
--	tolyfenpyrad	METI insecticide	Mitochondrial complex I electron transport inhibitor	21A

Rimon Spray Timing

- Rimon is best used early season as it only affects Lygus nymphs and there is more synchronization of the Lygus generations at that time
- Later application is best when tank mixed with another product
- Timing is critical



Contact Poison - Little Residual Activity

Pesticide	Chemical	Subgroup	Target Site of Activity	IRAC #
Malathion	malathion	organophosphate	acetylcholine esterase inhibitor	1B
Dibrom	naled	organophosphate	acetylcholine esterase inhibitor	1B
Brigade	bifenthrin	pyrethroid	sodium channel modulator	3
Danitol	fenpropathrin	pyrethroid	sodium channel modulator	3
Actara	thiamethoxam	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Assail	acetamiprid	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Rimon	novaluron	benzoylureas	inhibitor of chitin biosynthesis, type 0	15
Prevam	borax	borax	nonspecific inhibitor	uncl
Belay	clothianidin	neonicotinoid	nicotinic Ach receptor agonist/ antagonist	4A
Beleaf	flonicamid	flonicamid	nonspecific feeding blocker	9C
Closer	sulfoxaflor	sulfilimine	--	uncl
--	tolyfenpyrad	METI insecticide	Mitochondrial complex I electron transport inhibitor	21A

May be useful in a mixture

Unregistered Products

Pesticide	Chemical	Subgroup	Target Site of Activity	IRAC #
Malathion	malathion	organophosphate	acetylcholine esterase inhibitor	1B
Dibrom	naled	organophosphate	acetylcholine esterase inhibitor	1B
Brigade	bifenthrin	pyrethroid	sodium channel modulator	3
Danitol	fenpropathrin	pyrethroid	sodium channel modulator	3
Actara	thiamethoxam	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Assail	acetamiprid	neonicotinoid	nicotinic Ach receptor agonists/ antagonist	4A
Rimon	novaluron	benzoylureas	inhibitor of chitin biosynthesis, type 0	15
Prevam	borax	borax	nonspecific inhibitor	uncl
Belay	clothianidin	neonicotinoid	nicotinic Ach receptor agonist/ antagonist	4A
Beleaf	flonicamid	flonicamid	nonspecific feeding blocker	9C
Closer	sulfoxaflor	sulfilimine	--	uncl
--	tolyfenpyrad	METI insecticide	Mitochondrial complex I electron transport inhibitor	21A

- Nerve poison
- Feeding blocker
- Affects insect's energy metabolism
- ??

Lygus Control - Insecticides - 2011

Treatment	Chemical name	Rate (form/ac)
Untreated		
Brigade WSB	bifenthrin	16 oz
Actara 25 WG + Brigade	thiamethoxam + bifenthrin	4 oz + 16 oz
Rimon 0.83 EC	novaluron	12 oz
Rimon + Brigade	novaluron + bifenthrin	12 oz + 16 oz
Beleaf 50SG	flonicamid	2.8 oz
Belay 2.13 EC	clothianidin	4.0 oz
Belay 2.13 EC	clothianidin	6.0 oz
tolfenpyrad 15 SC	tolfenpyrad	27 oz
tolfenpyrad 15 EC	tolfenpyrad	27 oz
sulfoxaflor (L)	sulfoxaflor	2.85 oz
sulfoxaflor (M)	sulfoxaflor	4.28 oz
sulfoxaflor (H)	sulfoxaflor	5.7 oz

Treatments applied September 30, 2011, to first year 'Albion' field
All include Dyne-amic

Lygus Control - Small Nymphs - 2011

Treatment	Mean \pm SE small Lygus bug nymphs per plant							
	10/6/11		10/13/11		10/20/11		10/27/11	
Untreated	0.08	\pm 0.01	0.16	\pm 0.06	0.06	\pm 0.04	0.04	\pm 0.01
Brigade	0.02	\pm 0.02*	0.06	\pm 0.02	0.02	\pm 0.02	0.08	\pm 0.04
Actara + Brigade	0.03	\pm 0.02	0.10	\pm 0.06	0.08	\pm 0.04	0.05	\pm 0.04
Rimon	0.01	\pm 0.01*	0.06	\pm 0.02	0.05	\pm 0.02	0.11	\pm 0.04
Rimon + Brigade	0.03	\pm 0.01*	0.04	\pm 0.03	0.05	\pm 0.01	0.06	\pm 0.02
Beleaf	0.03	\pm 0.01*	0.03	\pm 0.00*	0.04	\pm 0.02	0.09	\pm 0.05
Belay (L)	0.03	\pm 0.01*	0.03	\pm 0.01*	0.02	\pm 0.02	0.02	\pm 0.02
Belay (H)	0.02	\pm 0.01*	0.01	\pm 0.01*	0.03	\pm 0.01	0.09	\pm 0.02
tolfenpyrad 15 SC	0.02	\pm 0.01*	0.03	\pm 0.03*	0.01	\pm 0.01	0.08	\pm 0.04
tolfenpyrad 15 EC	0.01	\pm 0.01*	0.06	\pm 0.03	0.05	\pm 0.01	0.12	\pm 0.03
sulfoxaflor (L)	0.00	\pm 0.00*	0.04	\pm 0.01	0.05	\pm 0.04	0.06	\pm 0.04
sulfoxaflor (M)	0.02	\pm 0.01*	0.04	\pm 0.01	0.01	\pm 0.01	0.05	\pm 0.01
sulfoxaflor (H)	0.05	\pm 0.01	0.08	\pm 0.02	0.04	\pm 0.01	0.04	\pm 0.02

* Means are significantly different from control at $P < 0.05$ using Dunnett's test.

Lygus Control - Total Lygus - 2011

Treatment	Mean \pm SE Total Lygus bugs (adults + nymphs) per plant							
	10/6/11		10/13/11		10/20/11		10/27/11	
Untreated	0.11	\pm 0.01	0.21	\pm 0.06	0.17	\pm 0.05	0.23	\pm 0.05
Brigade	0.08	\pm 0.02	0.13	\pm 0.04	0.11	\pm 0.03	0.23	\pm 0.03
Actara + Brigade	0.05	\pm 0.03	0.15	\pm 0.06	0.11	\pm 0.04	0.14	\pm 0.03
Rimon	0.06	\pm 0.02	0.09	\pm 0.02	0.08	\pm 0.02	0.19	\pm 0.02
Rimon + Brigade	0.04	\pm 0.02	0.06	\pm 0.05	0.10	\pm 0.04	0.10	\pm 0.03
Beleaf	0.08	\pm 0.02	0.06	\pm 0.01	0.09	\pm 0.02	0.17	\pm 0.07
Belay (L)	0.04	\pm 0.03	0.06	\pm 0.01	0.03	\pm 0.01	0.07	\pm 0.02
Belay (H)	0.02	\pm 0.01	0.01	\pm 0.01	0.12	\pm 0.04	0.12	\pm 0.03
tolfenpyrad 15 SC	0.09	\pm 0.03	0.07	\pm 0.07	0.06	\pm 0.05	0.11	\pm 0.07
tolfenpyrad 15 EC	0.03	\pm 0.01	0.10	\pm 0.04	0.09	\pm 0.02	0.19	\pm 0.04
sulfoxaflor (L)	0.06	\pm 0.03	0.10	\pm 0.02	0.09	\pm 0.06	0.15	\pm 0.06
sulfoxaflor (M)	0.02	\pm 0.01	0.11	\pm 0.04	0.07	\pm 0.03	0.16	\pm 0.03
sulfoxaflor (H)	0.08	\pm 0.01	0.12	\pm 0.02	0.10	\pm 0.03	0.09	\pm 0.03

Pre-treat count = 0.061 small Lygus nymphs per plant, 0.022 large Lygus nymphs per plant, and 0.015 Lygus adults per plant, for a total of 0.098 Lygus per plant

Percent fruit damage at 3, 4 and 5 weeks

Treatment	Percent fruit damaged per plot					
	10/20/11		10/27/11		11/3/2011	
Untreated	16.09	± 1.22	8.64	± 2.84	24.35	± 7.33
Brigade	11.66	± 1.03	9.85	± 3.94	9.57	± 4.91
Actara + Brigade	9.79	± 0.25	8.45	± 0.17	11.55	± 2.78
Agri-flex	11.55	± 2.06	7.16	± 2.55	9.90	± 2.80
Rimon	7.77	± 1.29	4.24	± 1.34	9.31	± 1.19
Rimon + Brigade	5.17	± 0.78	4.19	± 0.97	7.23	± 0.33
Beleaf	10.73	± 2.28	8.99	± 1.06	12.83	± 1.30
Belay (L)	9.59	± 3.04	5.78	± 2.52	6.31	± 2.80
Belay (H)	13.17	± 2.16	4.91	± 1.32	11.36	± 3.99
tolfenpyrad 15 SC	10.74	± 1.10	7.03	± 2.60	11.73	± 2.40
tolfenpyrad 15 EC	12.08	± 1.97	4.76	± 1.88	8.16	± 0.91
sulfoxaflor (L)	12.37	± 2.86	7.42	± 1.86	7.06	± 2.16
sulfoxaflor (M)	10.57	2.04	7.77	1.79	10.03	3.00
sulfoxaflor (H)	10.53	± 2.28	7.59	± 1.21	10.25	± 2.72

Percent fruit damage reduction at 3, 4 and 5 weeks

Treatment	Percent damage reduction			
	10/20/11	10/27/11	11/3/11	Average
Untreated				
Brigade	27.52	(14.11)	52.57	21.99
Actara + Brigade	39.16	2.10	59.33	33.53
Agri-flex	28.21	17.08	61.76	35.68
Rimon	51.72	50.90	70.30	57.64
Rimon + Brigade	67.87	51.44	47.30	55.54
Beleaf	33.34	(4.15)	74.08	34.42
Belay (L)	40.39	33.02	53.34	42.25
Belay (H)	18.14	43.18	51.83	37.72
tolfenpyrad 15 SC	33.26	18.55	66.50	39.44
tolfenpyrad 15 EC	24.93	44.88	70.99	46.93
sulfoxaflor (L)	23.13	14.11	58.81	32.02
sulfoxaflor (M)	34.30	9.98	57.88	34.05
sulfoxaflor (H)	34.55	12.10	60.69	35.78

Western Flower Thrips

Frankliniella occidentalis



California



Type I Bronzing

Bronzing

3 types identified

Type I



Type III



Type II



Causes of Type 3 Bronzing



Koike, S.T., F.G. Zalom, and K.D. Larson. 2009. Bronzing of strawberry fruit as affected by production practices, environmental factors, and thrips. *HortScience*. 44(6): 1-6.

Causes of Type 3 Bronzing

Elevated temperature and solar radiation

Mitigated by:

overhead sprinkling

certain foliar pesticides

lignin

Koike, S.T., F.G. Zalom, and K.D. Larson. 2009. Bronzing of strawberry fruit as affected by production practices, environmental factors, and thrips. HortScience. 44(6): 1-6.


Western Flower Thrips Control, Orange Co.

Treatment	Number of thrips per flower			
	Feb 18	Feb 27	Mar 4	Mar 16
Untreated	1.14 ±0.62	5.29 ±1.94	6.90 ±2.72	11.10 ±3.52
Lannate	0.47 ±0.46	0.73 ±0.35*	1.87 ±1.16*	6.87 ±3.10
Entrust	0.45 ±0.30	1.39 ±0.45*	2.98 ±0.81*	6.15 ±0.89*
Success	0.60 ±0.72	1.58 ±0.86*	3.70 ±2.29	7.87 ±2.14

* Treatment differs from untreated by pairwise t-test at $P < 0.05$.

- Problem - concern for resistance and restriction on number of applications per season by Dow Agrosiences for all spinosyns (Entrust, Success and Radiant) on strawberries in the Monterey Bay area.
- Other registered insecticides are not very effective.

Thrips Susceptibility of Fields, Pretreatment



Thrips insecticide efficacy - Watsonville, 2011

Treatment	Rate (form/ac)	Mean \pm SE thrips adults per flower		
		10/7/11 ¹	10/13/11 ²	10/27/11 ³
Untreated	NA	1.06 \pm 0.24	0.61 \pm 0.36	1.22 \pm 0.47
Actara + Brigade	4.0 oz + 16 oz	1.11 \pm 0.24	1.83 \pm 0.38	0.94 \pm 0.15
Tolfenpyrad 15SC	27 fl oz	0.39 \pm 0.39	0.78 \pm 0.46	1.22 \pm 0.20
Tolfenpyrad 15EC	27 fl oz	0.28 \pm 0.20	1.33 \pm 0.33	0.94 \pm 0.45
Closer	2.85 fl oz	1.28 \pm 0.34	1.06 \pm 0.24	1.44 \pm 0.45
Closer	4.28 fl oz	1.28 \pm 0.28	1.11 \pm 0.87	1.61 \pm 0.59
Closer	5.7 fl oz	0.78 \pm 0.34	1.28 \pm 0.45	1.33 \pm 0.38

Tolfenpyrad and Closer are not registered for strawberries

Corn Earworm

Helicoverpa zea



UC Statewide IPM Project
© 2000 Regents, University of California



UC Statewide IPM Project
© 2000 Regents, University of California



Project
University of California

Corn Earworm Control, Orange County, 2008

Trade name	Chemical name	Rate form./acre
Untreated	--	--
Altacor	chlorantraniliprole	3.0 oz
Altacor	chlorantraniliprole	6.0 oz
Rimon	novaluron	12.0 oz
Intrepid	methoxyfenozide	10.0 oz
Synapse	flubendiamide	2.0 oz
Synapse	flubendiamide	3.0 oz
Radiant SG	spinetoram	8.0 oz

Application date 4/11/2008

Corn Earworm Control, Orange County, 2008

Treatment	Rate (form/A)	<i>H zea</i> damaged fruit per 20 plants
Untreated		3.31 ± 0.69
Altacor	3.0 oz	1.08 ± 0.21*
Altacor	6.0 oz	1.28 ± 0.39*
Rimon	12.0 fl oz	1.72 ± 0.35*
Intrepid	10.0 fl oz	1.90 ± 0.23*
Synapse WG	2.0 oz	1.60 ± 0.58*
Synapse WG	3.0 oz	1.23 ± 0.64*
Radiant SG	8.0 fl oz	1.90 ± 0.47*

Application date 4/11/2008

Harvest date 4/22/2008

ANOVA statistics - $F=6.4435$, $df=7,23$ $P=0.001$

Corn Earworm Control, Watsonville, 2011

Treatment	chemical	Rate (form/acre)	Mean \pm SE damaged fruit per 16 plants
Untreated	--	NA	5.25 \pm 1.38a
DiPel DF ¹	Bt	0.25 lb	1.75 \pm 0.48b
DiPel DF ¹	Bt	0.50 lb	1.25 \pm 0.63b
DiPel DF ¹	Bt	1.0 lb	2.50 \pm 0.29b
Entrust	spinosad	1.5 oz	1.50 \pm 0.96b

- Means followed by the same letter do not differ significantly by Student's t-test at $P=0.05$.

¹ Applied in 40 gal. per acre

Application date 6/29/2011

Harvest date 9/6/2011

Corn Earworm Control, Watsonville, 2011

Treatment	chemical	Rate (form/acre)	Mean \pm SE damaged fruit per 20 plants
Untreated	--	--	6.5 \pm 0.5a
Altacor	chlorantraniprole	4.5 oz	1.3 \pm 0.6b
Synapse	flubendiamide	4.5 oz	1.5 \pm 0.4b
Intrepid	methoxyfenozide	10 oz	2.0 \pm 0.5b
Rimon	novaluron	12 oz	1.8 \pm 0.4b
Radiant	spinetoram	10 oz	2.0 \pm 1.2b

* Means followed by the same letter do not differ significantly by Student's t-test at $P=0.05$.

Application date(s) 8/16/2011 and 8/23/2012 (Bt only)

Harvest date 7/19/2011

ANOVA statistics - $F=8.9714$, $df=5,23$ $P=0.0002$

Vinegar flies - *Drosophila* spp.

An phytosanitary export and processing concern

Quality assurance protocol for export to Japan developed in 1998

Cultural controls

More frequent harvests

Sanitation



Vinegar flies - *Drosophila* spp.

Camarillo, 2012



Vinegar flies - *Drosophila* spp.

Sources



Vinegar flies - *Drosophila* spp.

Traps will capture many species and other flies, too

Traps contain either
baker's yeast + sugar + water
or apple cider vinegar



Vinegar flies - *Drosophila* spp.

Insecticides won't control the maggots once fruit are infested, they can only knock down adult flies and protect uninfested fruit.

Organophosphates

Malathion

Pyrethroids*

Danitol, (Brigade, Bifenture)

Spinosyns

Entrust, Success, Radiant

* *Using pyrethroids can exacerbate Lygus resistance*

Spotted Wing Drosophila

Drosophila suzukii

New species in North America
Attacks sound fruit
Problem for fresh market



Drosophila melanogaster
and other species

Always present
Attacks older fruit
Problem for processing



Spotted Wing Drosophila

Damage to strawberry



Spotted Wing Drosophila

Identification and Biology




Male



Female

Adults are 2-3 mm in size. Females and their larvae (maggots) are easily confused with other *Drosophila*

Spotted Wing *Drosophila*



Other *Drosophila*



Drosophila suzukii

D. suzukii has a specialized sharp ovipositor, different from other *Drosophila*.

Spotted Wing Drosophila

Management is same as for all *Drosophila*

- Sanitation, remove mature and overripe fruit
- Sanitation, eliminate alternate habitat (culled fruit, abandoned host fields) that sustains the infestation
- Monitoring and trapping to quickly detect infestations - get ahead of the damage
- Use insecticidal sprays or baits to suppress fly populations
 - organophosphates, pyrethroids, spinosyns

*11th Annual Strawberry Production Meeting in Ventura County
Camarillo, August 31, 2012*

Lygus, Thrips, Corn Earworm and Spotted Wing Drosophila

Frank Zalom

Department of Entomology
University of California, Davis